

Nova[®] 40W

EPA Reg. No. 62719-411

**Section 18 Quarantine Exemption for
Control of Rust (*Phakopsora pachyrhizi*) in Beans and Peas
(Crop Groups 6, 7)[†]****(For Use and Distribution Only in the State of North Dakota)****ATTENTION**

- **Section 18 Quarantine Exemption:** This use is authorized under EPA quarantine exemption pursuant to Section 18 of the Federal Insecticide Fungicide and Rodenticide Act as amended.
- **This quarantine exemption is effective upon USDA confirmation that soybean rust is present in the continental U.S. and expires May 31, 2009.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Nova[®] 40W fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Nova 40W according to this Section 18 labeling is subject to all use precautions and limitations imposed by the label affixed to the product container for Nova 40W.

Directions for Use

[†]Including, but not limited to: Asparagus Bean, Bean, Blackeyed Pea, Broad Bean (Succulent), Chinese Longbean, Cowpea, Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, Garden Pea, Green Pea, Jackbean, Lima Bean (Green), Moth Bean, Pea, Pigeon Pea, Runner Bean, Snap Bean, Snow Pea, Southern Pea, Soybean (Immature Seed), Sugar Snap Pea, Sword Bean, Wax Bean, Yardlong Bean, and plant parts of any legume vegetable that will be used as animal feed

Apply 4 to 5 oz of Nova 40W (1.6 to 2 oz ai myclobutanil) per acre, using ground or aerial equipment, in an adequate spray volume to achieve good coverage and canopy penetration. For aerial application, apply Nova 40W in a minimum spray volume of 10 gallons per acre. Begin applications when rust is first observed. Make subsequent applications at 7- to 10-day intervals if conditions remain favorable for disease development.

Environmental Hazards

Groundwater: This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface water: This chemical can contaminate surface water through spray drift. Under some conditions, this chemical may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

Specific Use Restrictions:

- **Do not** allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.
- **Do not** feed bean and pea forage and hay to livestock.

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- **Do not** make more than 2 applications of myclobutanil-containing products to beans and peas per season.
- **Do not** apply more than 0.625 lb of Nova 40W (0.25 lb ai myclobutanil) per acre per crop.
- **Rotation Crops** – Fields treated with myclobutanil (Laredo[®], Nova and Rally[®]) can be rotated at any time to crops that are listed on a registered myclobutanil label immediately after the last treatment. Do not plant other crops within 30 days after the last application of a product containing myclobutanil.

Any adverse effects resulting from the use of Nova 40W under this quarantine exemption must be immediately reported to the North Dakota Department of Agriculture.

[®]Trademark of Dow AgroSciences LLC

R866-034

File Code: 06-ND-09

Accepted: 07/31/06

Initial printing.